

Exploring trans-Planckian physics and the curvature effect by primordial power spectrum with WMAP five-year data

Jie Ren¹, Hong-Guang Zhang², and Xin-He Meng²

¹*Theoretical Physics Division, Chern Institute of Mathematics, Nankai University, Tianjin 300071, China and*

²*Department of physics, Nankai University, Tianjin 300071, China*

(Dated: March 11, 2009)

The vacuum inflation with a boundary condition specified at a short-distance scale in a generally primeval non-flat Universe, and implications of the correspondingly modified primordial power spectrum via the WMAP data are investigated. We obtain a general form of the modified primordial power spectrum including the effects of both possibly new physics scale and the spatial curvature. The modulation of the primordial power spectrum due to new physics is of the order H/Λ , where H is the Hubble parameter during inflation and Λ is the new physics scale that is key to inflation, while the modulation from the curvature effect is of the order K/k^2 , where K is the spatial curvature before inflation and k is the comoving wave number. We also find that a closed Universe before inflation is favored by WMAP five-year data.

PACS numbers: 98.80.Cq, 98.70.Vc

I. INTRODUCTION

Great progress achieved in observational cosmology in the past few years has offered us an opportunity to probe the possible trans-Planckian physics as one of high energy frontiers in the spectrum of cosmic microwave background (CMB) radiation fluctuations. Inflation is increasingly convinced to be a consistent picture for universe evolution by mainly CMB experiments [1]. The physics at high energy scales is reflected in the cosmic size magnified by microscopic quantum fluctuations during the inflation era and the trans-Planckian physics in inflationary cosmology has been studied and proposed to be probably detectable [2, 3, 4, 5, 6]. In astrophysics observations the WMAP data have been analyzed to constrain cosmological models and study physics during inflation with the primordial power spectrum [7, 8, 9]. The imprints of quantum fluctuations implied by a quantum gravity theory near Planck scale, like superstring theory is potentially detectable with more precise data like in the upcoming Planck satellite mission, which will help us to explore the fundamental laws of nature.

Many aspects of new physics have been studied recently, partly such as the CPT violation [10], temperature of the inflation [11], modified dispersion relation [12], noncommutative space time [13], modified uncertainty principle [14], relations to dark energy [15] and various other interesting topics. A simple but well motivated case is that, for an energy scale Λ , new physics emerges as a boundary condition in the momentum space without assuming the specific nature of the short-distance physics. The effect can be described by a modulation term to the primordial power spectrum. Some versions of the modulation of the primordial power spectrum present as a linear term [2, 3], while others show that the leading order is quadratic [16, 17]. Some authors claim that the effects are linear in H/Λ is of crucial importance to have detectable effects [4] since $H \sim 10^{14}$ GeV during the inflation. In literature the non-flat pre-inflation Universe

has been studied [18, 19] and the modified primordial power spectrum due to the spatial curvature has been calculated out [18], as another perspective to explore the trans-Planckian physics.

We perform a combined analysis of the modulation to the primordial power spectrum due to both the new physics scale and the spatial curvature. Following Danielsson's method [2], we calculate the exact solution of the primordial power spectrum, and then propose an approximate form with explicit physical meanings. The leading order modulation to the primordial power spectrum by new physics is a term with order of magnitude H/Λ , where H is the Hubble parameter during inflation and Λ is the new physics scale, important for inflation. And the leading order modulation by the curvature effect is a K/k^2 term, where K is the spatial curvature before inflation and k is the comoving wave number. Based on this theoretical result, we can add two more parameters, which describe the new physics scale and the curvature contributions, into the six-parameter standard model of cosmology and employ the WMAP five-year data to fit globally the parameters.

This paper is organized as follows: In the next section, we present the basic calculations of the primordial power spectrum. In Sec III, we summarize the method and the results obtained by Danielsson and Easter *et al.* Then we combine the effects of the possibly new physics scale and the spatial curvature into the primordial power spectrum. In Sec. IV, we use the newly modified primordial power spectrum to fit the WMAP data sets. Finally, we present our conclusions with discussions in the last section.

II. BASICS OF THE FRAMEWORK

First considering the Friedmann-Robertson-Walker metric in the flat space geometry ($K=0$) as the case fa-

vored by observational data

$$ds^2 = -dt^2 + a(t)^2 d\mathbf{x}^2, \quad (1)$$

where $a(t)$ is the scale factor. The Hubble parameter is defined by $H = \dot{a}/a$, where the dot denotes a derivative with respect to comoving time t . In conformal coordinates the metric takes the form

$$ds^2 = a(\eta)^2 (-d\eta^2 + d\mathbf{x}^2), \quad (2)$$

where η is the conformal time. We take $\eta = -\frac{1}{aH}$ as an approximation if H is near constant during inflation. By solving Einstein's equation, we obtain the Friedmann equations

$$\frac{3}{a^2} \left(\frac{a'}{a} \right)^2 = \frac{8\pi}{M_{\text{p}}^2} \rho, \quad -\frac{1}{a^2} \left[\frac{2a''}{a} - \left(\frac{a'}{a} \right)^2 \right] = \frac{8\pi}{M_{\text{p}}^2} p, \quad (3)$$

where ρ and p are respectively the globally isotropic and homogenous energy density and the pressure for the cosmological fluid assumed as perfect fluid, and a prime denotes a derivative with respect to conformal time. The content of the early Universe is assumed to be described by a scalar field ϕ with a potential $V(\phi)$ from a fundamental theory, and the equation of state (EOS) can be expressed as $\rho = \frac{1}{2}\dot{\phi}^2 + V(\phi)$, with $p = \frac{1}{2}\dot{\phi}^2 - V(\phi)$. The equation of energy conservation $\dot{\rho} + 3H(\rho + p) = 0$ for the scalar field is $\ddot{\phi} + 3H\dot{\phi} + V'(\phi) = 0$.

The simple inflationary scenario predicts that a single scalar field generates adiabatic, Gaussian and nearly scale invariant spectra of scalar and tensor perturbations. We use the rescale field μ that is determined by an equation of parametric oscillation

$$\mu''_{\text{S,T}} + (k^2 - \frac{z''_{\text{S,T}}}{z_{\text{S,T}}})\mu_{\text{S,T}} = 0, \quad (4)$$

where $z_{\text{S}} = \sqrt{\epsilon}a$ and $z_{\text{T}} = a$. Here S and T denote the scalar and tensor perturbations, respectively, and $\sqrt{\epsilon} \simeq \dot{\phi}/H$ is the first slow roll parameter (we take $8\pi/M_{\text{p}}^2 \equiv 1$ for simplicity). For the detailed derivations, see Ref. [20]. The solution of Eq. (4) should satisfy the asymptotic condition

$$\lim_{k/(aH) \rightarrow \infty} \mu_{\text{S,T}}(\eta) = \frac{e^{-ik(\eta-\eta_i)}}{\sqrt{2k}}. \quad (5)$$

In this paper, we only consider the scalar perturbations and assume ϵ is almost constant. The primordial power spectrum for scalar perturbations is defined by

$$P(k) = \frac{k^3}{2\pi^2} \left| \frac{\mu}{\sqrt{\epsilon}a} \right|^2. \quad (6)$$

The scalar spectrum index and the running scalar spectrum index are defined respectively by

$$n_s - 1 \equiv \frac{d \ln P(k)}{d \ln k}, \quad \alpha \equiv \frac{d^2 \ln P(k)}{d(\ln k)^2}. \quad (7)$$

The solution of Friedmann equations combined with the EOS of perfect fluid is given by

$$a(\eta) \propto \eta^{2/(1+3w)}, \quad (8)$$

which can be written as $a(\eta) = \eta^{-c}$, if we write the EOS parameter as $w = -\frac{1}{3} - \frac{2}{3c}$ (which is obtained by a holographic dark energy model [21]). Note that both the power-law inflation $a(t) \propto t^p$ and the exponential inflation $a(t) \propto e^t$ in terms of comoving time correspond to a power-law form $a(\eta) \propto \eta^{-c}$ in terms of conformal time. The case $c = 1$, *i.e.*, $w = -1$ the cosmological constant by definition, is the case that we are most interested in, thus Eq. (4) is

$$\mu'' + \left[k^2 - \frac{c(c+1)}{\eta^2} \right] \mu = 0. \quad (9)$$

The exact solution of this equation can be explicitly given out as

$$u(\eta) = C_1 \sqrt{\eta} H_{c+1/2}(k\eta) + C_2 \sqrt{\eta} H_{c+1/2}^*(k\eta), \quad (10)$$

where the integral constants C_1 and C_2 are determined by the asymptotic condition Eq. (5), and $H_l(x)$ is the Hankel function. Since the asymptotic condition Eq. (5) describes a spherical wave in momentum space, we choose the spherical Hankel function $h_l(x)$. The solution should be $u(\eta) = C(k)\eta \cdot h_c^*(\eta k)$. In the following we can see that the effect of the new physics is included by a mixture of the two terms. Thus in the general case, the primordial power spectrum for scalar perturbations is given by

$$P(k) = \left(\frac{H}{\dot{\phi}} \right)^2 \left(\frac{H}{2\pi} \right)^2 \lim_{\eta \rightarrow 0} \eta^2 |C_1 h_c(k\eta) + C_2 h_c^*(k\eta)|^2. \quad (11)$$

The limit $\eta \rightarrow 0$ is because we consider the late times.

III. NEW PHYSICS SCALE AND THE CURVATURE EFFECT

We would like to discuss the encoded effects of trans-Planckian physics in the primordial power spectrum. The main procedure of a general approach proposed by Danielsson [2] is summarized as follows. The initial condition is imposed when $p = \Lambda$, where Λ is the energy scale of new physics important for inflation. We treat H/Λ as a free parameter and fit it with observational data in the next section. Since the comoving momentum is given by $k = ap = -\frac{\Lambda}{Hk}$, the initial condition is

$$\eta_0 = -\frac{\Lambda}{Hk}. \quad (12)$$

By defining the conjugate momentum

$$\pi_k = \mu'_k - \frac{a'}{a} \mu_k, \quad (13)$$

the quantization of the system in terms of time-dependent oscillators is given by

$$\mu_k(\eta) = \frac{1}{\sqrt{2k}}(a_k(\eta) + a_{-k}^\dagger(\eta)), \quad (14)$$

$$\pi_k(\eta) = -i\sqrt{\frac{k}{2}}(a_k(\eta) - a_{-k}^\dagger(\eta)). \quad (15)$$

Then we can apply a Bogoliubov transformation to mix the creation and annihilation operators at a specific conformal time η_0 . The choice of vacuum

$$a_k(\eta_0)|0, \eta_0\rangle = 0 \quad (16)$$

means that $v_k(\eta_0) = 0$ and $\pi_k(\eta_0) = ik\mu_k(\eta_0)$.

We choose the linear combination of the first-order spherical Hankel function and its conjugate

$$f_k = \frac{A_k}{\sqrt{2k}}e^{-ik\eta} \left(1 - \frac{i}{k\eta}\right) + \frac{B_k}{\sqrt{2k}}e^{ik\eta} \left(1 + \frac{i}{k\eta}\right), \quad (17)$$

as the solution of Eq. (9) in the case $c = 1$. The Wronskian condition

$$\mu_k^* \frac{du_k}{d\tau} - u_k \frac{d\mu_k^*}{d\tau} = -i \quad (18)$$

gives

$$|A_k|^2 - |B_k|^2 = 1. \quad (19)$$

The choice of vacuum $v_k(\eta_0) = 0$ presents

$$B_k = \frac{ie^{-2ik\eta_0}}{2k\eta_0 + i} A_k, \quad (20)$$

where $A_k \approx 1$ if η_0 is very large. A more general calculation for A_k and B_k modified by trans-Planckian physics can be found in Ref. [22]. Finally, the primordial power spectrum is obtained by

$$\begin{aligned} P(k) &= \frac{k^3}{2\pi^2 a^2} |f_k|^2 \\ &\sim \frac{1}{4\pi^2 \eta^2 a^2} (|A_k|^2 + |B_k|^2 - A_k^* B_k - A_k B_k^*) \\ &= \left(\frac{H}{2\pi}\right)^2 \left[1 - \frac{H}{\Lambda} \sin \frac{2\Lambda}{H} + \frac{H^2}{\Lambda^2} \sin^2 \frac{\Lambda}{H}\right]. \end{aligned} \quad (21)$$

This result is for the gravitational wave, and one needs to take an extra factor $1/\epsilon$ into account for scalar perturbations. Easter *et al.* generalize this result to an arbitrary background [3]

$$u_k = \frac{1}{2} \sqrt{\frac{\pi}{k}} \sqrt{\frac{y}{1-\epsilon}} \left[C_+ H_\nu \left(\frac{y}{1-\epsilon} \right) + C_- H_\nu^* \left(\frac{y}{1-\epsilon} \right) \right]. \quad (22)$$

Their result can be written as

$$P(k) = \left[1 + \frac{H}{\Lambda} \sin \frac{2\Lambda}{(1-\epsilon)H} + \frac{H^2}{\Lambda^2} \sin^2 \frac{2\Lambda}{(1-\epsilon)H} \right] P_0(k), \quad (23)$$

where $P_0(k)$ denotes the primordial power spectrum in Bunch-Davies vacuum. This is consistent with Danielsson's result, since the sign before the modulation term is insignificant.

We further generalize the modulation of the primordial power spectrum to non-flat Universe. The effects of the spatial curvature before inflation have been studied in Ref. [18]. The calculations for the open Universe and the closed Universe are slightly different, thus we only show the closed Universe for example. For the detailed calculations, please see Ref. [18]. Note that a non-zero curvature is assumed before inflation, while the present curvature is assumed to be zero as universe large expansion. In a closed Friedmann Universe, the scale factor for a closed Universe can be solved as

$$a(t) = \frac{\sqrt{K}}{H} \cos Ht, \quad (24)$$

where K is the curvature of the space. The scale factor in terms of the conformal time is

$$a(\eta) = -\frac{\sqrt{K}}{H} \frac{1}{\sin \sqrt{K}\eta}, \quad (25)$$

where $\frac{\sqrt{K}}{H} < a < \infty$ as $-\frac{\pi}{2\sqrt{K}} < \eta < 0$. Then, the perturbation equation is

$$\mu_k'' + \left[k^2 - K \left(2 \csc^2(\sqrt{K}\eta) + 1 \right) \right] \mu_k = 0. \quad (26)$$

The general solution of this equation is given by

$$\begin{aligned} \mu_k &= C_1 \left[\sqrt{k^2 + K} - i\sqrt{K} \cot(\sqrt{K}\eta) \right] e^{-i\sqrt{k^2 + K}\eta} \\ &+ C_2 \left[\sqrt{k^2 + K} + i\sqrt{K} \cot(\sqrt{K}\eta) \right] e^{i\sqrt{k^2 + K}\eta}, \end{aligned} \quad (27)$$

where

$$C_1 = \frac{A_k}{\sqrt{2}(k^2 + K)^{3/4}}, \quad C_2 = \frac{B_k}{\sqrt{2}(k^2 + K)^{3/4}}. \quad (28)$$

We use Eq. (11) to calculate the modulation of the primordial power spectrum by spatial curvature as

$$P(k) = \frac{1}{(1 + K/k^2)^{3/2}} P_0(k). \quad (29)$$

In the following we follow Danielsson's method to encode the new physics scale that is important for inflation into the primordial power spectrum for a universe with non-zero curvature. It is easy to check that the Wronskian condition also gives $|A_k|^2 - |B_k|^2 = 1$. The choice of vacuum gives

$$B_k = \frac{k - \sqrt{k^2 + K} + i\sqrt{K} \cot(\sqrt{K}\eta_0)}{k + \sqrt{k^2 + K} + i\sqrt{K} \cot(\sqrt{K}\eta_0)} e^{-2i\sqrt{k^2 + K}\eta_0} A_k. \quad (30)$$

We consider the case at late times when $\eta \rightarrow 0$ similar to Eq. (21), and obtain the primordial power spectrum as

$$P(k) = \left[1 + \frac{(k - \sqrt{k^2 + K})^2 + K \cos(2\sqrt{k^2 + K}\eta_0)}{2k\sqrt{k^2 + K}} - \frac{\sqrt{K} \cot(\sqrt{K}\eta_0) \sin(2\sqrt{k^2 + K}\eta_0)}{k} + \frac{K \cot^2(\sqrt{K}\eta_0) \sin^2(\sqrt{k^2 + K}\eta_0)}{2k\sqrt{k^2 + K}} \right] \times \frac{1}{(1 + K/k^2)^{3/2}} P_0(k), \quad (31)$$

which is the final result with the modulation of the primordial power spectrum in a generally closed Universe. This can be reduced to Eqs. (21) and (29).

By assuming that corrections from both new physics and the curvature effect are very small, we can use the linear approximation, in which, the modified primordial power spectrum is given by

$$P(k) = A_s k^{-2\epsilon} \left[1 - \xi k^{-\epsilon} \sin\left(\frac{2}{\xi k^{-\epsilon}}\right) - \frac{K}{k^2} \right], \quad (32)$$

where the possibly new physics gives a $\xi = H/\Lambda$ factor correction to the primordial power spectrum, and the spatial curvature contributes a K/k^2 term, before which the coefficient can be absorbed in K . For the open Universe with negative K , we can also obtain the same form as Eq. (32). The physical meaning is clear: If no new physics, the asymptotic behavior of the spherical Hankel function is the free spherical wave. The cutoff due to new physics serves as a boundary condition in momentum space, and therefore, the choice of the vacuum leads to a mixture of the positive and negative mode states, *i.e.*, outward- and inward-going waves classically. The amplitude of the spherical wave in a non-flat space will be modified and contribute a separate term by linear approximation. The modulations of the primordial power spectrum will be magnified in the CMBR fluctuations, which can be probed by WMAP mission and the future PLANCK satellite observations.

IV. PROBING NEW PHYSICS AND THE CURVATURE EFFECTS BY WMAP5 DATA

Reconstructing the primordial power spectrum [23, 24, 25] from CMB experiments is an essential task to understand the early Universe physics. The commonly assumed form of the primordial power spectrum is

$$P(k) = A_s \left(\frac{k}{k_0} \right)^{n_s - 1}, \quad (33)$$

where $n_s = 1$ corresponds to the Harrison-Zeldovich (scale invariant) power spectrum. Slow-roll inflationary models predict that n_s is very close to one. To study

more details for the spectrum, we should also concern on the second-order term. A standard approach is the logarithmic expansion to $P(k)$ as below

$$\ln P(k) = \ln A_s + (n_s - 1) \ln \left(\frac{k}{k_0} \right) + \frac{\alpha}{2} \ln^2 \left(\frac{k}{k_0} \right). \quad (34)$$

The modification of the power spectrum due to new physics can be highly model-dependent, nevertheless, we can make as less as possible assumptions in our model. The modified primordial power spectrum can be tested with more and more observational CMB data with increasingly precision. As we know Danielsson's result of the modification is

$$P(k; \epsilon, \xi) = P_0(k) \left\{ 1 - \xi \left(\frac{k}{k_n} \right)^{-\epsilon} \sin \left[\frac{2}{\xi} \left(\frac{k}{k_n} \right)^{\epsilon} \right] \right\} \quad (35)$$

where $\xi = 4 \times 10^{-4} \sqrt{\epsilon}/\gamma$, and $\gamma = \Lambda/M_p$, and the constant k_n corresponds to the largest scales measurable in the CMB [7]. As another approach, the oscillations in the primordial power spectrum was studied by setting a step in the inflationary potential [26]. In the present work, we focus on the curvature effect in the data fitting.

We propose a general modified primordial power spectrum in form as

$$P(k; \epsilon, \xi, K) = A_s \left(\frac{k}{k_0} \right)^{-2\epsilon} (1 - M_\xi - M_K), \quad (36)$$

where

$$M_\xi = \xi \left(\frac{k}{k_0} \right)^{-\epsilon} \sin \left[\frac{2}{\xi} \left(\frac{k}{k_0} \right)^{\epsilon} \right], \quad M_K = \frac{K'}{(k/k_0)^2}. \quad (37)$$

Here M_ξ denotes the modulation due to new physics, and M_K stands for the modulation from the spatial curvature. We make a data fitting analysis to the obtained modified primordial power spectrum by using Monte Carlo techniques as implemented in the publicly available CosmoMC code [27] together with the likelihood code developed by the WMAP team [1]. We take $k_0 = 0.05 \text{ Mpc}^{-1}$.

The parameters in our model are the six parameters in the standard cosmology model plus two addition parameters that describe the new physics scale and the spatial curvature, respectively. We have numerically calculated the scalar spectrum index n_s and its running α as a function of the wave number k , *i.e.*, $n_s = n_s(k)$ and $\alpha = \alpha(k)$. Since k serves as a parameter, we cannot obtain the exact values of n_s and α , but we can plot the relation between them. Fig. 1 plots the relation between n_s and α , which is consistent with the values fitted by the WMAP team. We set $\xi = 0$ and fit the model with data, thus only K' is a new parameter in the model. Firstly we does restrict the value of K' to be positive or negative, but after the fitting we find that the closed Universe with positive K' is favored. Fig. 2 plots the marginalized contours of the




FIG. 1: The relation between the scalar spectrum index and its running. Here we take $K' = 10^{-6}$ and the solid, dashed, and the dotted lines correspond to the parameters (ϵ, ξ) as $(0.024, 0.015)$, $(0.024, 0.01)$, $(0.02, 0.015)$, respectively.




FIG. 2: The marginalized contours (68% and 95% C.L.) for the model only with curvature.

parameters n_s and K' . If we take $K' \sim 10^{-6}$, the energy density of the spatial curvature today is

$$\Omega_K \sim \frac{K' k_0^2 c^2}{a_0^2 H_0^2} \sim 10^{-2}. \quad (38)$$

In conclusion, the result shows that a model for the primordial power spectrum with a positive curvature is slightly favored by WMAP five-year data fittings.

V. CONCLUSION AND DISCUSSION

We have obtained a general modulation formula to the CMB primordial power spectrum due to both possibly new physics and the curvature effect. The leading order of additional terms in the modified primordial power spectrum is H/Λ and K/k^2 , respectively. From the classical point of view, the effects of new physics can be regarded as a reflection result by the boundary of the momentum space, *i.e.*, a mixture of the outward- and inward-going waves. The spatial curvature before inflation should not be too large, otherwise it will largely change the observed power spectrums, and thus a fine-tuning may be required. Nevertheless, it is reasonable to include the curvature generally and indeed it improves the data fitting. We find that a closed pre-inflation Universe is slightly favored by the data fittings.

Another important issue in cosmology is to understand the nature of the dark energy dominated current cosmic speed-up expansion. It may be probable that the inflation and the dark energy can be described by a unified theory, which exhibits different dynamic and kinematic behaviors in the early high energy scale universe and the later low energy scale universe evolutions. Some approaches such as modified gravity [28] has been attempted to explore such possibilities. The observational cosmology has challenged the situation that the string theory is believed as beyond the experimental science, but cosmic observations will provide powerful data sets to test the implications from high energy scale quantum gravity, like string cosmology [29]. We can be sure that the upcoming astrophysics missions, like Planck satellite will open a new era for us to understand the underlying physics in inflation scenario and dark energy mystery, with expectations to conceive possibly new physics and fundamental laws of nature.

ACKNOWLEDGEMENT

We acknowledge the use of the Legacy Archive for Microwave Background Data Analysis (LAMBDA). Support for LAMBDA is provided by the NASA Office of Space Science. J.R. thanks Prof. Richard Easther for helpful comments. X.H.M. thanks Prof. David Lyth for helpful discussions. X.H.M. was supported by NSFC under No.10675062. This work was supported by Nankai university ISC.

[1] C.L. Bennett *et al.*, *Astrophys. J. Suppl.* **148**, 1 (2003), astro-ph/0302207; D.N. Spergel *et al.*, astro-ph/0603449; E. Komatsu, *et al.*, arXiv:0803.0547 [astro-ph].

[2] U.H. Danielsson, *Phys. Rev. D* **66**, 023511 (2002).

[3] R. Easther, B.R. Greene, W.H. Kinney, and G. Shiu, *Phys. Rev. D* **66**, 023518 (2002).

- [4] L. Bergstrom and U.H. Danielsson, *JHEP* **0212**, 038 (2002).
- [5] R. Easther, B.R. Greene, W.H. Kinney, and G. Shiu, *Phys. Rev. D* **64**, 103502 (2001); **67**, 063508 (2003).
- [6] C.P. Burgess, J.M. Cline, F. Lemieux, and R. Holman, *JHEP* **0302** 048 (2003).
- [7] O. Elgaroy and S. Hannestad, *Phys. Rev. D* **68**, 123513 (2003).
- [8] R. Easther, W.H. Kinney, and H. Peiris, *JCAP* **0505**, 009 (2005).
- [9] J. Martin and C. Ringeval, *JCAP* **0608**, 009 (2006).
- [10] B. Feng, M. Li, J.Q. Xia, X. Chen, and X. Zhang, *Phys. Rev. Lett.* **96**, 221302 (2006).
- [11] K. Bhattacharya, S. Mohanty, and R. Rangarajan, *Phys. Rev. Lett.* **96**, 121302 (2006).
- [12] R.H. Brandenberger and J. Martin, *Mod. Phys. Lett. A* **16**, 999 (2001); J. Martin and R.H. Brandenberger, *Phys. Rev. D* **63**, 123501 (2001); *Phys. Rev. D* **65**, 103514 (2002).
- [13] C.S. Chu, B.R. Greene, and G. Shiu, *Mod. Phys. Lett. A* **16**, 2231 (2001); R. Brandenberger and P.M. Ho, *Phys. Rev. D* **66**, 023517 (2002); Q.G. Huang and M. Li, *Nucl. Phys. B* **713**, 219 (2005); X. Zhang and F.Q. Wu, *Phys. Lett. B* **638**, 396 (2006).
- [14] S.F. Hassan and M.S. Sloth, *Nucl. Phys. B* **674**, 434 (2003).
- [15] L. Mersini, M. Bastero-Gil, and P. Kanti, *Phys. Rev. D* **64**, 043508 (2001); B. Chen, M. Li, and Y. Wang, *astro-ph/0611623*.
- [16] A. Kempf and J.C. Niemeyer, *Phys. Rev. D* **64**, 103501 (2001).
- [17] N. Kaloper, M. Kleban, A. Lawrence, and S. Shenker, *Phys. Rev. D* **66**, 123510 (2002).
- [18] E. Masso, S. Mohanty, and G. Zsembinski, *astro-ph/0609349*.
- [19] G. Efstathiou, *Mon. Not. Roy. Astron. Soc.* **343**, L95 (2003); A. Lasenby and C. Doran, *Phys. Rev. D* **71**, 063502 (2005); D.H. Lyth and E.D. Stewart, *Phys. Lett. B* **252**, 336 (1990); D.H. Lyth and A. Woszczyna, *Phys. Rev. D* **52**, 3338 (1995); B. Allen, R.R. Caldwell, and S. Koranda, *Phys. Rev. D* **51**, 1553 (1995); A.A. Starobinsky, *astro-ph/9603075*; M. Bucher, A.S. Goldhaber, and N. Turok, *Phys. Rev. D* **52**, 3314 (1995); S. Gratton, T. Hertog, and N. Turok, *Phys. Rev. D* **62** (2000) 063501.
- [20] J. Martin and D.J. Schwarz, *Phys. Rev. D* **57**, 3302 (1998).
- [21] M. Li, *Phys. Lett. B* **603**, 1 (2004).
- [22] A. A. Starobinsky, *JETP Lett.* **73**, 371 (2001), *astro-ph/0104043*.
- [23] J.E. Lidsey, A.R. Liddle, E.W. Kolb, E.J. Copeland, T. Barreiro, and M. Abney, *Rev. Mod. Phys.* **69**, 373 (1997).
- [24] D.H. Lyth and A. Riotto, *Phys. Rept.* **314**, 1 (1999).
- [25] S. Hannestad, *JCAP* **0404**, 002 (2004).
- [26] J. Hamann, L. Covi, A. Melchiorri, and A. Slosar, *Phys. Rev. D* **76**, 023503 (2007).
- [27] A. Lewis and S. Bridle, *Phys. Rev. D* **66**, 103511 (2002).
- [28] S. Nojiri and S.D. Odintsov, *Phys. Rev. D* **68**, 123512 (2003); S. Capozziello, S. Nojiri, and S.D. Odintsov, *Phys. Lett. B* **632**, 597 (2006); X.H. Meng and P. Wang, *Phys. Lett. B* **584**, 1 (2004); *Class. Quant. Grav.* **22**, 23 (2005); **21**, 2029 (2004); **21**, 2527 (2004).
- [29] U.H. Danielsson, *Class. Quant. Grav.* **22**, S1, (2005).